


Fundamental Physics with Black Hole Binaries

K. Chatziioannou (Flatiron)
A. Klein (U of Birmingham)
H. Witek (King's College London)

Florence, November 14 2018

BHBs in the LISA band



(LISA proposal '17)

- 10s - 100s MBHBs/yr (Klein+ '16)
- 10s - 100s SOBHBs/yr (Sesana '16)

Massive BHs in GR

Massive BHBs in GR are (or will soon be) including a range of effects:

- Accurate inspiral phasing
- Multiple waveform harmonics
- Spin-induced precession
- Orbital eccentricity
- Merger signal
- Multiple ringdown harmonics
- Kicks

All can be affected by GR modifications

Massive black holes – beyond standard model

Gravity

- Is general relativity correct?
 - Test in new regimes
- Black holes and light fields
- Black holes in modified gravity
 - Exotic compact objects

Cosmology

- What is dark matter?
- What is dark energy?

High-energy physics


- Quantum gravity?
- BSM particles?

smoking guns:


- additional radiation → dephasing of GW signal, new effects?
- additional polarizations, ringdown/echoes, ...
- inconsistency with GR prediction of full IMR – or is it?

Methods – overview

- gravitational wave detections are **theory-driven**
- waveform templates needed for **identification** and **interpretation** of signal
- so far: (almost) only in GR, i.e., we are deaf to any “non-vanilla” models!
- **waveform-free** analyses are also being developed (data analysis session)



FYI: LISA Waveform WG



(LISA Management Plan '18)

- consider joining LISA Waveform Working group
(co-chairs: D. Shoemaker, M. van de Meent, N. Warburton, H. Witek)
- wav-wg@lisamission.org

Waveforms in GR: current status

PN/Phenom:

- IMR, full precession, subdominant harmonics, precession+harmonics, small eccentricity
- inspiral, eccentricity, full precession, subdominant harmonics

EOB:

- IMR, eccentricity, full precession, subdominant harmonics

Surrogate:

- Based on NR: moderate spins, circular, moderate mass ratio, full mode content (= what NR has)

Inspiral modifications: parametrized

Phasing and amplitude modifications: parametrized post-Einsteinian description

(Yunes & Pretorius '09)

Modification of the Fourier domain phase: $\Psi(f) = \sum_n a_{\text{GR}} f^{n/3} + \beta f^{b/3}$.

Modification of the Fourier domain amplitude: $A(f) = A_{\text{GR}}(f)(1 + \alpha f^{a/3})$

LIGO implementation: TIGER (Agathos+ '14)

- Modular
- Can be adapted to different WF models
- e.g. dynamical scalarization: $\Psi(f) = \sum_n a_{\text{GR}} f^{n/3} + \theta(f - f_*)\beta f^{b/3}$
- Spin precession?
- Eccentricity?
- Number of parameters grow for each new effect included


Inspiral++ modifications: theory-specific

- Scalar-tensor, PN (Damour & Esposito-Farèse '96, Mirshekari & Will '13, Lang '14, ...)
- Scalar-tensor, EOB (Julié & Deruelle '17)
- Einstein-Maxwell-dilaton, EOB (Khalil+ '18)
- Avoid exponential growth in number of parameters
- Have to adapt each WF model for each alternative theory and search for modifications separately

Numerical Relativity in GR I

State-of-the-art (see, e.g., review by Duez & Zlochower '18)


- performed regularly for
 - comparable-mass $q \lesssim 1/15$
 - moderate spins
 - quasi-circular binaries
- large spins
(Lovelace et al '08, '11; Scheel et al '14;
Ruchlin, Healy, Lousto, Zlochower '16, 17; Hinder et al '18)
- waveform catalogues
(e.g. SXS/SpEC, RIT, GAT, ...)




(SXS/SpEC '13)

Practical timescales: $\mathcal{O}(1\text{month})$ per simulation

Numerical Relativity in GR II



(Chu et al '16)



(Nakano et al '11)


New challenges

- better accuracy
- low-intermediate mass ratios $q \lesssim 1/100$
(Lousto et al '10; Nakano et al '11; Sperhake et al '11)
- eccentric binaries
- Longer duration; more precession cycles
- (accurate) higher harmonics; inform ringdown studies (excitation factors)
- triple systems? Environmental effects?

Numerical relativity beyond GR I

- Black holes and light fields (Okawa, Witek, Cardoso '14; Zilhão, Witek, Cardoso '15; Pretorius & East '17, East '17, '18)
 - so far: only single BHs \rightarrow monochromatic GWs
 - (M)BHB signatures? resonances & hang-ups? Dephasing?
 - so far: only weak-field approximation



- exotic compact objects
 - (scalar) boson stars (Liebling & Palenzuela '12, Palenzuela et al '17, Helfer et al '18, Bezares et al '18, ...),
 - Proca stars (Sanchis-Gual et al '18)



Example: boson stars (Helfer et al '18)

upcoming: CGQ Focus Issue “Numerical relativity beyond GR”, eds. D. Gerosa, L. Stein, H. Witek


Numerical relativity beyond GR II



Example: sGB (Witek, Gualtieri, Pani, Sotiriou '18)

- BHs in modified gravity
 - compact binaries in scalar-tensor theory
(Barausse et al '12, Shibata et al '13, Healy et al '11, Berti et al '13)
 - Einstein-Maxwell-Dilaton models
(Hirschmann et al '17)
 - dynamical Chern-Simons gravity
(Okounkova et al '17)
 - scalar Gauss-Bonnet gravity
(Witek et al '18)
- relevance for LISA vs LIGO/3G; MBHBs vs SOBHBs?
- “full” models vs EFT-type approaches; well-posed IVPs

Ringdown modifications




(Berti+ '16)

Parametrized approach: post-Kerr formalism (Glampedakis+ '17)

Link to specific theories?

Waveform accuracy requirements

Comprehensive accuracy requirement calculations are necessary




(Mangiagli et al '18)

- Fairly insensitive to the mass ratio
- Mostly depends on the total mass/time to coalescence

Waveform accuracy requirements

Unsurprisingly, a GR systematic (here lack of higher modes) can mimic beyond-GR effects (here ppE-type merger-inspiral terms)



(Pang et al. arxiv:1802.03306)

Testing GR – current status

A number of currently operational tests of GR

- Parametric deviations (very waveform dependent)
- Residual power (waveform dependent)
- Consistency test (less waveform dependent)
- Propagation effects (very waveform dependent)
- Polarization tests (less waveform dependent)

Parametric deviations – very waveform dependent


$$\tilde{h} \sim \tilde{A}(f; \vec{\theta}_{GR}) e^{i \sum_i p_i(\vec{\theta}_{GR}) f^i}$$

$$p_i \rightarrow p_i(1 + \delta p_i)$$

Theory	a	α	b	β
Brans-Dicke	–	0	-7/3	β
Parity-Violation	1	α	0	–
Variable $G(t)$	-8/3	α	-13/3	β
Massive Graviton	–	0	-1	β
Quadratic Curvature	–	0	-1/3	β
Extra Dimensions	–	0	-13/3	β
Dynamical Chern-Simons	+3	α	+4/3	β

Yunes+ (arxiv:0909.3328)

Cornish+ (arxiv:1105.2088)



LVC (arxiv:1706.01812)


- Can be repeated for any BBH
- Relies heavily on an accurate background GR waveform

Propagation effects – very waveform dependent

$$E^2 = p^2 c^2 + A p^\alpha c^\alpha$$

$$\delta v_g \sim \frac{\text{GW period}}{\text{travel time}} \sim \frac{\text{GW wavelength}}{\text{distance}}$$

For 800Mpc and 250Hz, $\delta v_g \sim 5 \times 10^{-20}$



LVC (arxiv:1706.01812)


$$\delta v_g \sim A E^{\alpha-2} \Rightarrow A \sim \delta v_g E^{2-\alpha} \sim \delta v_g (h_{\text{Pl}} f)^{2-\alpha} \sim 10^{-20} \text{peV}^{2-\alpha}$$

For 2Gpc and 10^{-3}Hz , $\delta v_g \sim 5 \times 10^{-15}$ and $A \sim 10^{-15} \text{aeV}^{2-\alpha}$


- Can be repeated for any BBH
- Relies heavily on an accurate background GR waveform

Residual power – waveform dependent

Subtract a (=maxL, maxP, fair draw) template waveform and check for residual power



LVC (arxiv:1602.03837)




LVC (arxiv:1602.03841)


Cornish+ (arxiv:1410.3835)

- Can be repeated for any BBH
- Can capture generic unmodeled deviations
- Relies on an accurate template to subtract
- Perhaps limited applicability if many overlapping sources (data analysis session)

Consistency test – less waveform dependent




LVC (arxiv:1602.03841)



LVC (arxiv:1706.01812)

- Can be repeated for any BBH with a ringdown
- Relies on an accurate mapping between initial and final masses and spins
- In principle better ringdown tests with MBH (access to more ringdown harmonics)

Testing GR – more under construction



Will ([arxiv:1403.7377](https://arxiv.org/abs/1403.7377))

- Continuous waves from bosonic clouds (Isi, Sun, Brito, Melatos '18)
- Null-stream test for beyond-GR polarization content (Guersel, Tinto '89)
- Parametrized test for beyond-GR polarization content (Chatziioannou, Yunes, Cornish '12)
- Modes consistency (Dhanpal, Ghosh, Mehta, Ajith, Sathyaprakash '12)
- Multipolar consistency (Kastha, Gupta, Arun, Sathyaprakash, van den Broeck+ '18)
- Theory-specific tests
- etc.

Discussion

- What (curvature/energy) regimes can we test with LISA vs LIGO / 3G?

Discussion

- Which beyond-GR models are more informative to study?
- Toy models vs fundamental models? How informative are they?

Discussion – “We need more NR simulations”

- Needed for synergy with PN/EOB and perturbation theory
- Next directions? What is feasible?
- Fill parameter space vs informed model building
- What are physically interesting / well-motivated theories to simulate?
- good news: “EFT-type” (\sim parametrized) approach adaptable to many models

Discussion – “We need better waveforms”

- Depends on effect in mind (parameter hierarchy)
- Accuracy for parametrized tests vs residuals vs global fits
- What do we need to prioritize (inspiral, merger, ringdown, eccentricity, precession, mass ratio...?)
- Astrophysical effects?
- Multibanding

Discussion – GW tests of GR

- Which tests are going to be the most useful?
- What are the current tests not probing?
- Null vs parametrized vs theory specific tests
- Consistency tests vs constraints
- How does the absence of beyond-GR effects help with theory development?